









MRT Results for S. PATIENT, page 1 of 2

Physician: **WELLNESS, MARK MD**
 Patient: **SAMPLE PATIENT**
 Identifier: **S70123**
 Profile: **MRT Test ML150**
 Test Date: **07/10/2012**
 Technician: **EH**



Oxford Clinical Laboratory
 3555 Fiscal Court, Suite #9
 Riviera Beach, FL 33404

Phone: (561) 848-7111
 FL License #: L800010492
 CLIA ID #: 10D0914874
 U.S. Patents: 6,114,174 6,200,815



MRT Results for S. PATIENT, page 2 of 2

Physician: **WELLNESS, MARK MD**
 Patient: **SAMPLE PATIENT**
 Identifier: **S70123**
 Profile: **MRT Test ML150**
 Test Date: **07/10/2012**
 Technician: **EH**



Oxford Clinical Laboratory
 3555 Fiscal Court, Suite #9
 Riviera Beach, FL 33404

Phone: (561) 848-7111
 FL License #: L800010492
 CLIA ID #: 10D0914874
 U.S. Patents: 6,114,174 6,200,815

Meats & Poultry	Reaction Level
EGG	Non-Reactive
LAMB	Non-Reactive
BEEF	Non-Reactive
TURKEY	Non-Reactive
CHICKEN	Non-Reactive
PORK	Non-Reactive

Miscellaneous	Reaction Level
COLA	Non-Reactive
TAPIOCA	Non-Reactive
COFFEE	Non-Reactive
BAKER'S YEAST	Non-Reactive
HOPS	Reactive
TEA	Reactive

Beans/Nuts/Legumes	Reaction Level
LENTIL	Non-Reactive
ALMOND	Non-Reactive
SUNFLOWER SEED	Non-Reactive
HAZELNUT	Non-Reactive
PECAN	Non-Reactive
GREEN PEA	Non-Reactive
SOYBEAN	Non-Reactive
PISTACHIO	Non-Reactive
WALNUT	Non-Reactive
GARBANZO BEAN	Non-Reactive
CASHEW	Non-Reactive
PINTO BEAN	Moderately Reactive
LIMA BEAN	Moderately Reactive
PEANUT	Moderately Reactive

Degrees of reactivity may not in all cases correlate with levels of clinical reactivity to the tested food or chemical. It is recommended to completely avoid all tested items with "Reactive," or "Moderately Reactive" scores. If tested "Non-Reactive" foods have been consumed regularly before testing, there is a high probability that they are "safe" and are not likely to provoke symptoms, as long as there is no history of allergic, autoimmune, or other type of inflammation-provoking or symptom-provoking reaction.

These test results can form the basis of an eating plan developed under the guidance of a physician, dietitian, or other qualified healthcare practitioner.